

Math 123: Linear D.E.s of First and Second Order

Ryan Blair

CSU Long Beach

Tuesday October 11, 2016

Outline

- 1 First Order Differential Equations
- 2 Integrating Factor Method
- 3 Superposition Principle

Types of Differential equations

Definition

A first order **linear** D.E. is of the form

$$y' + Q(x)y = R(x)$$

where $Q(x)$ and $R(x)$ are functions of x .

Types of Differential equations

Definition

A first order **linear** D.E. is of the form

$$y' + Q(x)y = R(x)$$

where $Q(x)$ and $R(x)$ are functions of x .

Definition

A second order **linear** D.E. is of the form

$$y'' + P(x)y' + Q(x)y = R(x)$$

where $P(x)$, $Q(x)$ and $R(x)$ are functions of x .

Types of Differential equations

Definition

A first order **linear** D.E. is of the form

$$y' + Q(x)y = R(x)$$

where $Q(x)$ and $R(x)$ are functions of x .

Definition

A second order **linear** D.E. is of the form

$$y'' + P(x)y' + Q(x)y = R(x)$$

where $P(x)$, $Q(x)$ and $R(x)$ are functions of x .

If $R(x) = 0$ we call the D.E. **homogeneous**.

Integrating Factor Method

Question 1: Given a D.E. $y' + Q(x)y = R(x)$, if you could find a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

could you solve the D.E.?

Integrating Factor Method

Question 1: Given a D.E. $y' + Q(x)y = R(x)$, if you could find a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

could you solve the D.E.?

Answer: Yes!

Integrating Factor Method

Question 1: Given a D.E. $y' + Q(x)y = R(x)$, if you could find a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

could you solve the D.E.?

Answer: Yes!

Question 2: Given a D.E. $y' + Q(x)y = R(x)$, can you find the formula for a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

Integrating Factor Method

Question 1: Given a D.E. $y' + Q(x)y = R(x)$, if you could find a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

could you solve the D.E.?

Answer: Yes!

Question 2: Given a D.E. $y' + Q(x)y = R(x)$, can you find the formula for a function $f(x)$ such that

$$\frac{d(f(x)y)}{dx} = f(x)(y' + Q(x)y)$$

Answer: Yes! $f(x) = e^{\int Q(x)dx}$

Integrating Factor Method

To solve $y' + Q(x)y = R(x)$,

- 1 Multiply both sides by $f(x) = e^{\int Q(x)dx}$
- 2 Recognize that the L.H.S. is $\frac{d(f(x)y)}{dx}$
- 3 Integrate both sides and solve for y .

Exercise: Solve $y' + 2y = 2e^x$.

Exercise: Solve $xy' + y = \sqrt{x}$.

Superposition Principle

Theorem

Given a homogeneous linear differential equation with solutions $f(x)$ and $g(x)$ then $a \cdot f(x) + b \cdot g(x)$ is also a solution for any constants a and b .

Exercise: Demonstrate this theorem for the D.E.

$$y'' + P(x)y' + Q(x)y = 0$$